

# Dynamics of imidacloprid resistance and cross-resistance in the brown planthopper, *Nilaparvata lugens*

Yan Hua Wang<sup>1,2†</sup>, Sheng Gan Wu<sup>2†</sup>, Yu Cheng Zhu<sup>3\*</sup>, Jin Chen<sup>1</sup>, Feng Yi Liu<sup>4</sup>,  
Xue Ping Zhao<sup>2</sup>, Qiang Wang<sup>2</sup>, Zhen Li<sup>2</sup>, Xian Ping Bo<sup>1</sup> & Jin Liang Shen<sup>1\*</sup>

<sup>1</sup>Key Laboratory of Monitoring and Management of Plant Diseases and Insects, Ministry of Agriculture, Nanjing Agricultural University, Nanjing, Jiangsu 210095, China, <sup>2</sup>Institute of Quality and Standard for Agro-products, Zhejiang Academy of Agricultural Sciences, Hangzhou, Zhejiang 310021, China, <sup>3</sup>Jamie Whitten Delta States Research Center, ARS-USDA, Stoneville, MS 38776, USA, and <sup>4</sup>Huizhou Agricultural Technology Extension Center, Huizhou, Guangdong 516001, China

Accepted: 25 November 2008

**Key words:** resistance stability, Homoptera, Delphacidae, neonicotinoid insecticides, resistance management, rice

## Abstract

Extensive use of imidacloprid for suppressing the brown planthopper, *Nilaparvata lugens* (Stål) (Homoptera: Delphacidae), has placed heavy selection pressure on the target insect. A systematic study was carried out to determine imidacloprid resistance dynamics and cross-resistance. Data collected from a 3-year study (2005–2007) showed that in 2005, the resistance levels in Nanning (Guangxi), Haiyan (Zhejiang), and Nanjing and Tongzhou (Jiangsu) populations ranged from 200- to 799-fold compared with the susceptible strain. However, the resistance levels decreased to 135- to 233-fold in 2007, after reduced application of the chemical. A laboratory population was challenged with imidacloprid in successive generations. After 23 generations, the resistance ratio had increased from 200- to 1 298-fold. Continuous selection with imidacloprid could increase the resistance level even more than has already been developed in the population. Stopping selection with imidacloprid led to a rapid decrease of resistance from 759- to 114-fold after 17 generations. Resistance levels then became stable without decreasing any further. A similar result was also obtained from a study involving a field population (resistance ratio = 625-fold) collected from Tongzhou. At first, the population showed a rapid decrease in resistance right after imidacloprid selection was stopped, and then the resistance stabilized at a level of 105–129-fold. More interestingly, resistance increased again when selection was resumed. In addition, the resistant strain selected with imidacloprid showed substantial cross-resistance to imidaclothiz, thiacloprid, and acetamiprid, and slight levels of cross-resistance to dinotefuran and thiamethoxam, but no obvious cross-resistance to nitenpyram, buprofezin, and fipronil. The information from this study is valuable for formulating resistance-management strategies against *N. lugens*.

## Introduction

The brown planthopper, *Nilaparvata lugens* (Stål) (Homoptera: Delphacidae), is an economically important

insect of rice in Asia (Heinrich, 1994). Hopperburn, caused by severe feeding damage, is a serious threat to rice yields in most rice-growing areas (Ding & Su, 2002). Currently, chemical control is still a major method for suppressing *N. lugens* (Endo & Tsurumachi, 2001). Intensive use of imidacloprid has prompted adaptation of *N. lugens* to the chemical. Since the introduction of DDT, *N. lugens* has developed resistance to almost every insecticide introduced for *N. lugens* control. Each newly introduced chemical provided effective control of *N. lugens* at first, then became less effective years later. New chemicals have

\*Correspondence: Yu Cheng Zhu, USDA-ARS, Stoneville, MS 38776, USA. E-mail: yc.zhu@ars.usda.gov

Jin Liang Shen, Key Laboratory of Monitoring and Management of Plant Diseases and Insects, Ministry of Agriculture, Nanjing Agricultural University, Nanjing 210095, China.

E-mail: jlshen@njau.edu.cn

†Yan Hua Wang and Sheng Gan Wu contributed equally to this article.

been sought for better control of *N. lugens* because of these frequent chemical control failures (Kilin et al., 1981; Chung & Sun, 1983; Hirai, 1993).

Neonicotinoid insecticides, including imidacloprid, thiamethoxam, dinotefuran, nitenpyram, acetamiprid, thiacloprid, and imidaclothiz, are a well-established group of insecticides (Jeschke & Nauen, 2005). Many of these chemicals are still highly effective against most field populations of *N. lugens* (Tang et al., 2006). Imidacloprid was registered for controlling *N. lugens* on rice in the early 1990s. It quickly became the primary insecticide in many rice-growing areas in China because of its systemic nature and high efficacy against sucking insects (Sun et al., 1996). Continuous use of imidacloprid as a major insecticide for controlling *N. lugens* has resulted in a gradual decrease of efficacy against the pest. The application dose has had to be increased from 15 g a.i./ha in the 1990s to 60–120 g a.i./ha in 2005 in order to maintain effective control. However, in the Yangtze River Delta areas, such as Jiangsu, Zhejiang, Anhui, and other provinces, the control efficacy dropped from 95 to 60% even as application dose increased substantially (CY Diao, unpubl.). Although laboratory selection of *N. lugens* with imidacloprid has produced a highly resistant strain (Liu et al., 2003; Liu & Han, 2006), very little research has been done to survey and characterize imidacloprid resistance in field populations of *N. lugens*.

In an effort to understand the resistance mechanisms, researchers found that piperonyl butoxide synergized imidacloprid toxicity in green peach aphid [*Myzus persicae* (Sulzer)], cat flea [*Ctenocephalides felis* (Bouché)], house fly (*Musca domestica* L.), and tobacco whitefly [*Bemisia tabaci* (Gennadius)], suggesting that P450-mediated detoxification could be an important biochemical mechanism for imidacloprid resistance (Prabhaker et al., 1997; Wen & Scott, 1997; Richman et al., 1999; Choi et al., 2001; Nauen et al., 2002). Therefore, proper incorporation of these inhibitors with imidacloprid may help to abolish or suppress resistance in imidacloprid-resistant populations (Tang et al., 2006).

Cross-resistance is a potential risk that can limit the durability of an insecticide. Cross-resistance to various neonicotinoid insecticides was observed in field and laboratory-selected strains of *B. tabaci*, *Leptinotarsa decemlineata* (Say), and *Drosophila melanogaster* (Meigen) (Elbert & Nauen, 2000; Le Goff et al., 2003; Nauen & Denholm, 2005; Mota-Sanchez et al., 2006; Alyokhin et al., 2007). In these cases, reduced susceptibility to imidacloprid was linked to reduced susceptibility to other neonicotinoids, including thiamethoxam, acetamiprid, and nitenpyram. In *N. lugens*, cross-resistance to acetamiprid was also found in an imidacloprid-resistant strain developed in the laboratory (Liu et al., 2003). Therefore, research needs to be conducted to clarify whether

cross-resistance also exists between imidacloprid and other neonicotinoids.


Susceptibility recovery in resistant populations is important for resistance management (Lan & Zhao, 2001). Numerous studies have shown that resistance can become relatively stable once insects have developed significant resistance, and that declines in insecticide resistance could be slow after the insecticide is no longer used, as was observed in the resistance of *Rhipicephalus (Boophilus) microplus* (Canestrini) and *Carpocapsa pomonella* (L.) to DDT, and *M. domestica* to isolan (Stone, 1962; Barnes & Moffit, 1963; Georgiou, 1964). In contrast, in some cases such as pyrethroid resistance in *Helicoverpa armigera* (Hübner) and triazophos resistance in *Chilo suppressalis* (Walker) (Wu et al., 1996; Qu et al., 2005), resistance was not very stable and tended to decline after use of the chemical ceased.

*Nilaparvata lugens* has developed significant resistance to imidacloprid in China since 2005 (JL Shen, unpubl.), and other neonicotinoids as well as the non-neonicotinoids buprofezin and fipronil are currently being implemented in rice-growing areas for insect control and resistance management. However, it was not clear whether the imidacloprid-resistant populations had already developed cross-resistance to other neonicotinoids, and non-neonicotinoids such as buprofezin and fipronil, and whether stopping or reducing imidacloprid applications would allow susceptibility to be recovered in the resistant field populations. In order to develop better resistance management strategies, this study was designed to examine the change in imidacloprid resistance under different selection pressures, and to investigate whether cross-resistance to other neonicotinoids and to buprofezin and fipronil existed in the imidacloprid-resistant strain.

## Materials and methods

### Insecticides

Imidacloprid [95% technical product (TC)], acetamiprid (97% TC), and buprofezin (98.1% TC) were provided by Changlong Chemical Industrial Group (Changzhou, Jiangsu, China), Nitenpyram (95% TC) and imidaclothiz (95% TC) by Nantong Jiangshan Agrochemical (Nantong, Jiangsu, China), Thiamethoxam (97.2% TC) and dinotefuran [10% soluble concentrate (SL)] by Syngenta Investment (Shanghai, China), Thiacloprid (97.75% TC) by Tianjing Xingguang Chemical (Tianjing, China), and Fipronil (87% TC) by Bayer Cropscience Hangzhou (Hangzhou, Zhejiang, China). For the laboratory assays, emulsifiable concentrates were prepared by mixing each insecticide (technical grade) with 10% (wt/vol) Triton-X-100 and acetone.



**Figure 1** Map showing four locations in three provinces for collection of the brown planthopper, *Nilaparvata lugens*, from 2005 to 2007. 1, Nanning, Guangxi; 2, Haiyan, Zhejiang; 3, Nanjing, Jiangsu; and 4, Tongzhou, Jiangsu.

#### Local and temporal variation in imidacloprid susceptibility

To examine imidacloprid-resistance levels in different rice-growing areas, four populations of *N. lugens* were collected in August from 2005 to 2007 from Nanning (22°7'N, 108°3'E, Guangxi), Haiyan (30°5'N, 120°8'E, Zhejiang), Nanjing (32°0'N, 118°5'E, Jiangsu), and Tongzhou (32°1'N, 121°1'E, Jiangsu) in China (Figure 1). These locations were chosen based on the migration route of *N. lugens* from the southern into the northeastern part of China, the importance of rice production, and history of the insecticide applications in the areas. The field populations were collected from the same farms each year. Approximately 800 adults and 500–600 nymphs were collected from each site and were transferred to the greenhouse on the campus of Nanjing Agricultural University. The insecticide-free hybrid rice (Shanyou 63) at tillering to booting stage was used for maintaining insect colonies and subsequent bioassays. The field-collected insects were mass mated, and the third instars of  $F_1$  progenies were used for bioassays. All insect colonies were maintained at  $27 \pm 1^\circ\text{C}$  and L16:D8.

The bioassays were conducted by using the rice stem-dipping method (Zhuang et al., 1999) to examine dose-response of various populations to imidacloprid and other insecticides. Rice seedlings at tillering to booting stage were pulled out and washed thoroughly. Rice stems (about 10 cm in length) with roots were cut and air-dried to remove excess water. Three rice stems were grouped and

dipped into the appropriate insecticide test solution for 30 s. After the rice stems were air-dried for approximately 1 h, moistened cotton was used to wrap the rice roots. The treated rice stems were then placed in a 500-ml plastic cup. Twenty third instars of *N. lugens* were introduced into each plastic cup using a vacuum device. Each bioassay included 5–6 different concentrations of each chemical plus a distilled water-only control, and three plastic cups were arranged as replicates for each concentration. The treated insects were maintained at  $27 \pm 1^\circ\text{C}$  and L16:D8. Mortality was recorded after 96 h, except for the buprofezin treatment, which was recorded after 120 h due to its relatively slow action. The nymphs were considered dead if they failed to show movement after being gently prodded with a fine brush.

#### Resistance selection and resistance stability

The susceptible strain (S) of *N. lugens* was obtained from Zhejiang Chemical Industrial Group. This strain was originally collected in 1995 in a rice field near Hangzhou (Zhejiang, China), and the insects have been maintained on insecticide-free hybrid rice (Shanyou 63) for approximately 120 generations in the laboratory.

**Laboratory population (Lab).** A population of the second generation in 1993 was collected from a rice field near Nanjing (Jiangsu, China). The population was treated once every two generations with imidacloprid using a rice

seedling spray method (Wang et al., 1988) in the laboratory. After 2005, the Lab population was used as a starting population for imidacloprid resistance selection.

**Tongzhou population (TZ2006).** A population of the third generation in 2006 was collected from a rice field in Tongzhou (Jiangsu, China). This population was reared on insecticide-free rice and then used for evaluating susceptibility change in the population. The imidacloprid-resistant strain (R) was developed in laboratory from the Lab population after being selected with imidacloprid for an additional 23 generations.

The same rice stem-dipping method described above was adopted for resistance selection of the Lab population. Approximately 1000 third instars of every generation were treated with imidacloprid by the rice stem-dipping method and subsequently maintained at  $27 \pm 1^\circ\text{C}$  and L16:D8 for 4 days. Survivors were transferred to another cage containing fresh rice seedlings. The mortality for resistance selection was controlled to range between 40 and 70% in order to ensure sufficient survivors to develop and produce enough progeny for the subsequent insecticide selection (the treated concentration was similar to the  $\text{LC}_{50}$  value of imidacloprid against each generation of *N. lugens*).

To assess resistance stability to imidacloprid in *N. lugens*, the sixth generation of the Lab and TZ2006 populations (after being selected with imidacloprid) was reared on insecticide-free hybrid rice (Shanyou 63) for 17 and 16 generations, respectively, in the laboratory and  $\text{LC}_{50}$  values were determined every generation. To test resistance changes, resistance selection with imidacloprid was resumed with a part of the eighth generation of the TZ2006 population and  $\text{LC}_{50}$  values were determined every subsequent generation.

### Cross-resistance

The same rice stem-dipping method (Zhuang et al., 1999) was adopted to determine the toxicity of the neonicotinoids, buprofezin, and fipronil against the imidacloprid-resistant strain (R) and the Lab population. Cross-resistance in *N. lugens* was evaluated by examining a cross-resistance ratio for each compound of interest, which was calculated by dividing the  $\text{LC}_{50}$  of the imidacloprid-resistant R Lab strain by the  $\text{LC}_{50}$  imidacloprid reference Lab strain.


### Data analysis

Probit models of the Polo program (LeOra Software, 1997) were used to estimate parameters of dose-mortality regression for each rice stem-dipping bioassay. Mortality was corrected using Abbott's formula for each probit analysis (Abbott, 1925). The resistance ratio was calculated by dividing the  $\text{LC}_{50}$  value of a field population by the  $\text{LC}_{50}$  value of the susceptible strain, unless otherwise stated. Two  $\text{LC}_{50}$  values were considered to be significantly different if no overlap was found between their 95% confidence intervals. Mean tests were conducted using Student's t-test ( $P < 0.05$ ).


## Results

### Local and temporal variation of imidacloprid susceptibility

Data from the 3-year resistance monitoring study (2005–2007) revealed variable resistance ratios in the four populations (Figure 2). Among the four populations sampled in 2005, the Tongzhou population obtained the highest resistance ratio ( $\text{RR} = 799$ ), which was almost four times higher than the Nanning population, with the lowest resistance ratio of 200-fold. The other two populations (Nanjing and Haiyan) had resistance ratios of 558- and



**Figure 2** Differential resistance levels to imidacloprid in four populations of *Nilaparvata lugens*, surveyed in 2005, 2006, and 2007, compared with the susceptible strain. The error bars represent the 95% confidence intervals of the resistance ratios.



**Figure 3** Changes of imidacloprid resistance in the laboratory population (Lab) caused by different imidacloprid selection pressures, continuing selection, and cessation of selection. The error bars represent the 95% confidence intervals of the resistance ratios.

583-fold, respectively. In 2006, resistance ratios decreased in all four populations with the Nanjing population showing the largest (more than three times) drop compared to the previous year level. The Tongzhou population still maintained the highest resistance ratio, which was up to six times higher than the Nanning population. Local variations in 2007 decreased to a relatively lower level than those detected in the two previous years.

#### Resistance stability and changes


The Lab population of *N. lugens* was selected continuously (every generation) with imidacloprid for 23 generations in the laboratory (Figure 3). Resistance to imidacloprid in the Lab strain increased by 6.5-fold over the 23-generation selection period, that is, from 200-fold in the starting generation to 1298-fold in the 23rd generation. The results also indicated that the resistance ratio may not yet have approached a plateau or maximum, even though the insect had already developed 1298-fold resistance to imidacloprid. Continuing selection may further increase resistance in *N. lugens*. Meanwhile, to assess the stability of imidacloprid resistance in *N. lugens*, approximately 1000 third-instar nymphs of the sixth generation were used for a selection interruption test. A sharp decrease in resistance was detected immediately after removal of the selection pressure (Figure 3). After 17 generations without imidacloprid selection, the resistance ratio descended from 759- to 114-fold. During the last five generations tested, the resistance ratio fluctuated around 120-fold, suggesting that a slow or incomplete recovery of the susceptibility was expected once the population of *N. lugens* had developed resistance to the chemical.

Another experiment involved the Tongzhou population (TZ2006) for testing resistance stability. TZ2006 had developed 625-fold resistance to imidacloprid in the field. After being maintained on insecticide-free rice for 16 generations, resistance dropped to 105-fold (Figure 4). The first eight generations after selection was stopped showed the fastest rate of resistance loss. The resistance ratio of the ninth generation reached 122-fold, and only a slight change in resistance ( $RR = 105$ – $129$ -fold) occurred in the subsequent seven generations, suggesting that the population also reached a sustainable resistance level similar to that of the Lab strain. In addition, a portion of the eighth generation of the TZ2006 population was used to study resuming selection with imidacloprid. The resistance ratio in the population quickly increased from 155- to 678-fold after each of the eight subsequent generations were treated with imidacloprid, a figure very similar to the original resistance level (Figure 4).

#### Cross-resistance to other insecticides

Cross-resistance to eight other insecticides was determined by comparing the resistance ratio of a highly imidacloprid-resistant strain (R: selected from Lab with  $RR = 1298$ -fold) with the resistance ratio of a relatively low imidacloprid-resistant population (Lab:  $RR = 200$ -fold). The resistant strain (R) showed substantial cross-resistance to imidacloprid ( $t = 2.96$ , d.f. = 1,  $P < 0.05$ ), thiacloprid ( $t = 2.98$ , d.f. = 1,  $P < 0.05$ ), and acetamiprid ( $t = 1.73$ , d.f. = 1,  $P < 0.05$ ), and slight levels of cross-resistance to dinotefuran ( $t = 1.19$ , d.f. = 1,  $P > 0.05$ ) and thiamethoxam ( $t = 0.35$ , d.f. = 1,  $P > 0.05$ ). There was no cross-resistance to nitenpyram ( $t = 0.43$ , d.f. = 1,  $P > 0.05$ ), buprofezin ( $t = 0.6$ , d.f. = 1,  $P > 0.05$ ), and fipronil ( $t = 0$ , d.f. = 1,  $P > 0.05$ ) (Table 1).

**Figure 4** Changes of imidacloprid resistance in the Tongzhou population (TZ2006) influenced by different imidacloprid selection pressures, discontinuing selection, and resuming selection. The error bars represent the 95% confidence intervals of the resistance ratios.



## Discussion

Imidacloprid resistance might be a major factor contributing to the frequent outbreaks of *N. lugens* in recent years (Cheng et al., 2003; Cheng & Zhu, 2006; Gao et al., 2006; Wang & Wang, 2007). *Nilaparvata lugens* is a highly migratory insect and is able to travel long distances between the southern and the northeastern part of China (Cheng et al., 1979). Resistance development to insecticides in *N. lugens* was expected to be slow, because it was thought that resistance could be diluted in the process of migration (Zhuang et al., 2004).

However, imidacloprid has been an important chemical for controlling *N. lugens* in China from the early 1990s to 2005. The chemical was used to control *N. lugens* not only in the emigrating region but also in the immigrating region (Liu et al., 2003). Farmers applied imidacloprid to every generation of *N. lugens* to prevent its outbreak in many rice growing areas. Compared to imidacloprid, other insecticides were seldom used because of their relatively lower efficacy against *N. lugens*. The widespread and intensive use of imidacloprid has also been observed in other Asian countries (Wang & Wang, 2007). In addition, imidacloprid, being a systemic insecticide, exhibited prolonged residual activity, which is likely to generate persistent selection pressure for resistance development in *N. lugens* (Liu et al., 2003). Our laboratory selection demonstrated that *N. lugens* was able to achieve a very high level of resistance to imidacloprid under constant selection pressure. Further resistance surveys in 2005 indicated that four field populations of *N. lugens* developed very high resistance levels to imidacloprid (RR = 200–799-fold), and its efficacy against the insect decreased significantly in the

Yangtze River Delta areas, including Jiangsu, Zhejiang, Anhui, and other provinces in the same year (CY Diao, unpubl.). Therefore, widespread and intensive use of imidacloprid could be a major driving force for accelerating resistance development in *N. lugens*. Our 3-year resistance monitoring showed that all four populations had developed very high resistance levels to imidacloprid. Resistance development to certain insecticides due to their long application history can also be seen in other insects (Denholm et al., 1998; Mohan & Gujar, 2003; Sayyed et al., 2005; Zhao et al., 2006; Yu & McCord, 2007).

Because imidacloprid has been almost the only insecticide used for controlling *N. lugens* in most of the rice-growing area, migration no longer postpones resistance development substantially (Cheng & Zhu, 2006). The results from resistance monitoring indicated that resistance to imidacloprid in *N. lugens* substantially increased following the immigration of *N. lugens* from the southern into the northeastern part of China, that is, the resistance ratio increased from 200-fold in Nanning to 583-fold in Haiyan, and from 558-fold in Nanjing to 799-fold in the Tongzhou population in 2005. Similarly in 2006 and 2007, the resistance levels of the four populations fluctuated along the migration route following the same pattern as found in 2005. This phenomenon was closely associated with the widespread use of imidacloprid in southern China and other southern Asian countries. The northeastern-bound migration subsequently allowed highly resistant insects to move to the northeastern rice growing areas, where they then were subjected to further selection. Therefore, resistance to imidacloprid in *N. lugens* increased dramatically in the northeastern rice areas after migration.

**Table 1** Cross-resistance to various neonicotinoid insecticides and to buprofezin and fipronil in the imidacloprid-resistant strain of *Nilaparvata lugens*

Insecticide	Population	Slope (SE)	LC <sub>50</sub> (95% CI) mg a.i./l	$\chi^2$ (d.f.)	Resistance ratio <sup>1</sup>	Cross-resistance ratio <sup>2</sup>
Imidacloprid	S <sup>3</sup>	2.15 (0.12)	0.08 (0.05–0.11)	1.7 (3)	1.0	
	Lab	1.97 (0.18)	16.01 (13.36–19.39)	0.9 (2)	200.1	
	R	2.02 (0.16)	103.88 (86.07–128.79)	0.3 (3)	1 298.5	6.5
Imidaclothiz	S	2.09 (0.20)	0.33 (0.27–0.40)	0.7 (4)	1.0	
	Lab	2.47 (0.19)	15.22 (13.03–17.88)	2.6 (4)	46.1	
	R	3.04 (0.23)	27.09 (23.45–31.58)	2.9 (3)	82.1	1.8
Thiacloprid	S	1.35 (0.11)	13.50 (10.60–17.70)	1.2 (3)	1.0	
	Lab	2.57 (0.21)	25.27 (21.52–29.51)	4.5 (3)	1.9	
	R	3.09 (0.31)	46.48 (40.44–53.73)	4.7 (4)	3.4	1.8
Acetamiprid	S	2.46 (0.22)	7.55 (6.42–9.01)	1.0 (1)	1.0	
	Lab	2.21 (0.17)	8.27 (6.55–9.98)	2.6 (4)	1.1	
	R	2.13 (0.13)	12.86 (10.83–15.26)	3.2 (3)	1.7	1.6
Dinotefuran	S	2.71 (0.25)	0.14 (0.11–0.18)	0.7 (1)	1.0	
	Lab	2.31 (0.18)	0.19 (0.16–0.23)	4.1 (4)	1.4	
	R	2.48 (0.22)	0.27 (0.21–0.37)	2.8 (7)	1.9	1.4
Thiamethoxam	S	2.18 (0.16)	0.11 (0.088–0.12)	1.2 (2)	1.0	
	Lab	2.03 (0.19)	0.69 (0.58–0.83)	3.7 (4)	6.3	
	R	1.43 (0.17)	0.78 (0.55–1.14)	3.1 (4)	7.1	1.1
Nitenpyram	S	2.17 (0.21)	0.47 (0.25–0.61)	0.9 (2)	1.0	
	Lab	4.60 (0.36)	0.65 (0.58–0.73)	2.6 (4)	1.4	
	R	3.57 (0.23)	0.70 (0.61–0.79)	1.9 (2)	1.5	1.1
Buprofezin	S	1.25 (0.18)	0.066 (0.058–0.074)	1.7 (3)	1.0	
	Lab	2.65 (0.21)	0.21 (0.18–0.25)	2.4 (4)	3.4	
	R	2.38 (0.30)	0.24 (0.20–0.28)	2.1 (3)	3.6	1.1
Fipronil	S	2.15 (0.12)	0.039 (0.032–0.046)	2.9 (3)	1.0	
	Lab	2.48 (0.15)	0.12 (0.099–0.14)	3.1 (2)	3.1	
	R	2.84 (0.20)	0.12 (0.10–0.14)	2.9 (4)	3.1	1.0

<sup>1</sup>Resistance ratio: LC<sub>50</sub> value of R or Lab/LC<sub>50</sub> value of S.<sup>2</sup>Cross-resistance ratio: LC<sub>50</sub> value of R/LC<sub>50</sub> value of Lab.<sup>3</sup>S, susceptible strain to provide baseline to imidacloprid.

Compared to the resistance levels in 2005, the resistance ratios in all four populations dropped slightly in 2006 and 2007. Decreased imidacloprid use in 2006 and 2007 may have contributed to the decrease in resistance in *N. lugens*. In 2005, the regulation authority partially suspended imidacloprid use for *N. lugens* control in China. Buprofezin, fipronil, chlorpyrifos, isoprocarb, and dichlorvos were recommended as alternatives and as candidates for rotation. Therefore, the imidacloprid resistance level in *N. lugens* appears to have decreased due to the reduced selection by imidacloprid. This phenomenon suggests that rotating imidacloprid with other insecticides possessing no cross-resistance is an important practice for susceptibility recovery and for delaying resistance development to imidacloprid in *N. lugens*.

It is well known that resistance development in insects is the consequence of insecticide selection (Wang et al., 2005;

Amorim et al., 2007). Once the selection pressure is removed, resistance development will stop or even decline (Tabashnik et al., 1994; Wu et al., 1996). In this study, we observed that the resistance ratio of two populations decreased as expected when resistance selection was stopped. We further discovered that imidacloprid resistance in *N. lugens* rebounded immediately after the resumption of insecticide selection.

Once an *N. lugens* population has obtained a high level of resistance to imidacloprid, it might be able to maintain a sustainable level of resistance after removal of selection pressure for several generations. In this study, we found that resistance levels in both Tongzhou and in the Lab populations tended to decrease after the removal of selection pressure, and then stabilized at similar levels, that is, RR = 105–129-fold, for eight and five generations, respectively. However, complete recovery of susceptibility

was not seen, suggesting that a sustainable level of resistance was reached in the populations without insecticide selection. Similar situations have also been observed in other insects (Georghiou, 1964; Bauernfeind & Chapman, 1985; Zhao et al., 1993; Wu et al., 1996; Qu et al., 2005). In the field, resistance levels in *N. lugens* fluctuated, closely corresponding to the intensity of imidacloprid applications. The results of our 3-year field surveys were consistent with changes in resistance detected from laboratory selections.

Under laboratory conditions, the rate of resistance recession after removal of the selection pressure depended on genotypic composition and relative fitness of resistance genes in the colony (Georghiou, 1964, 1972). Although *N. lugens* was able to develop a high level of resistance to imidacloprid in the laboratory, resistance was not stable and declined after selection stopped. This phenomenon might be associated with an incompletely recessive trait of imidacloprid resistance controlled by multiple genes and significant fecundity and egg hatch costs (Zhao et al., 2000; Liu & Han, 2006; Baker et al., 2007). Achieving a homozygous colony was slow because a relatively low dose was used for the laboratory selection. After imidacloprid selection was stopped, heterozygous and homozygous susceptible individuals could quickly increase in frequency in the colony due to their fitness advantages over resistant individuals (Liu & Han, 2006; Baker et al., 2007). This phenomenon is consistent with the hypothesis that imidacloprid resistance is controlled by multiple incompletely recessive genes that have a fitness cost. Using alternative insecticides without cross-resistance to imidacloprid for *N. lugens* control could have relieved selection pressure on the target insect. Therefore, the resistance levels to imidacloprid declined in 2006 and 2007 in all four populations. On the other hand, alternating and rotating imidacloprid with other insecticides without cross-resistance may minimize or slow resistance development in the insect.

In addition to the direct examination of resistance development to imidacloprid in the target insect, risk assessment of cross-resistance is another important part of resistance management. Because cross-resistance might be involved in both functionally similar and dissimilar chemicals (Shen & Wu, 1995; Zhao et al., 1995; Elzen, 1997), assessment of cross-resistance must include not only neonicotinoid insecticides, but also non-neonicotinoids, which have the potential to be alternated or rotated with imidacloprid to control resistant populations of *N. lugens*. In this study, we detected substantial cross-resistance to imidacloprid, thiacloprid, and acetamiprid in a highly imidacloprid-resistant strain of *N. lugens*. The insects also showed slight levels of cross-resistance to dinotefuran and thiamethoxam, but not to nitenpyram. In addition, the

highly imidacloprid-resistant strain of *N. lugens* showed no cross-resistance to buprofezin (an insect growth regulator) and fipronil (a phenylpyrazole insecticide). Therefore, to relieve imidacloprid selection pressure and to recover imidacloprid susceptibility in *N. lugens*, we suggest that nitenpyram, buprofezin, and fipronil can be used in resistance management programs for alternation and rotation with imidacloprid.

## Acknowledgements

The authors appreciate Dr Andrew Li (USDA-ARS, Kerrville, TX, USA) and Dr Huarong Li (Iowa State University, Ames, IA, USA) for critical review of an early version of the manuscript. The study was partially supported by the research program of the Ministry of Agriculture in China for Replacing High Toxicity Pesticides, the Funding Program of Zhejiang Academy of Agricultural Sciences for New Scientists, and the Program of Resistance Monitoring and Management for the Rice Planthopper sponsored by the Chinese National Agro-tech Extension and Service Center in cooperation with Bayer Cropscience.

## References

- Abbott WS (1925) A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology* 18: 265–267.
- Alyokhin A, Dively G, Patterson M, Castaldo C, Rogers D et al. (2007) Resistance and cross-resistance to imidacloprid and thiamethoxam in the Colorado potato beetle *Leptinotarsa decemlineata*. *Pest Management Science* 63: 32–41.
- Amorim LB, Fontes de Oliveira CM, Rios FM, Regis L & Lobo Silva-Filha MHN (2007) Development of *Culex quinquefasciatus* resistance to *Bacillus sphaericus* strain IAB59 needs long term selection pressure. *Biological Control* 42: 155–160.
- Baker MB, Alyokhin A, Porter AH, Ferro DN, Dastur SR & Galal N (2007) Persistence and inheritance of costs of resistance to imidacloprid in Colorado potato beetle. *Journal of Economic Entomology* 100: 1871–1879.
- Barnes MMB & Moffit HR (1963) Resistance to DDT in the adult codling moth and reference curves for guthion and carbaryl. *Journal of Economic Entomology* 56: 722–725.
- Bauernfeind RJ & Chapman RK (1985) Nonstable parathion and endosulfan resistance in green peach aphids (Homoptera: Aphididae). *Journal of Economic Entomology* 78: 516–522.
- Cheng XN, Chen RC & Xi X (1979) Research on migration of brown planthopper, *Nilaparvata lugens* (Stål). *Acta Entomologica Sinica* 22: 1–21.
- Cheng XN, Wu JC & Ma F (2003) Brown Planthopper: Occurrence and Control, pp. 1–37. China Agricultural Press, Beijing, China.
- Cheng JA & Zhu ZY (2006) Analysis on the key factors causing the outbreak of brown planthopper in Yangtze Area, China in 2005. *Plant Protection* 32: 1–4.



- Choi BR, Lee SW & Yoo JK (2001) Resistance mechanisms of green peach aphid, *Myzus persicae* (Homoptera: Aphididae), to imidacloprid. *Korean Journal of Applied Entomology* 40: 265–271.
- Chung TC & Sun CN (1983) Malathion and MIPC resistance in *Nilaparvata lugens*. *Journal of Economic Entomology* 76: 1–5.
- Denholm I, Cahill M, Dennehy TJ & Horowitz AR (1998) Challenges with managing insecticide resistance in agricultural pests exemplified by the whitefly, *Bemisia tabaci*. *Philosophical Transactions of the Royal Society of London B Biological Sciences* 353: 1757–1767.
- Ding JH & Su JY (2002) *Agricultural Entomology*. China Agricultural Press, Beijing, China.
- Elbert A & Nauen R (2000) Resistance of *Bemisia tabaci* (Homoptera: Aleyrodidae) to insecticides in southern Spain with special reference to neonicotinoids. *Pest Management Science* 56: 60–64.
- Elzen GW (1997) Changes in resistance to insecticides in tobacco budworm populations in Mississippi, 1993–95. *Southwestern Entomologist* 22: 61–72.
- Endo S & Tsurumachi M (2001) Insecticide susceptibility of the brown planthopper and the white-backed planthopper collected from southeast Asia. *Journal of Pesticide Science* 26: 82–86.
- Gao XW, Peng LN & Liang DY (2006) Factors causing the outbreak of brown planthopper (BPH), *Nilaparvata lugens* Stål in China in 2005. *Plant Protection* 32: 23–25.
- Georghiou GP (1964) The stability of resistance to carbamate insecticides in the housefly after cessation of selection pressure. *Bulletin of the World Health Organization* 30: 85–90.
- Georghiou GP (1972) The evolution of resistance to pesticides. *Annual Review of Ecology and Systematics* 3: 133–168.
- Heinrichs EA (1994) Impact of insecticides on the resistance and resurgence of rice planthopper. *Planthoppers: Their Ecology and Management* (ed. by RF Denno & TJ Perfect), pp. 571–614. Chapman & Hall, New York, NY, USA.
- Hirai K (1993) Recent trends of insecticide susceptibility in the brown planthopper *Nilaparvata lugens* (Stål) (Homoptera: Delphacidae). *Applied Entomology and Zoology* 28: 339–346.
- Jeschke P & Nauen R (2005) Neonicotinoid insecticides. *Comprehensive Molecular Insect Science* (ed. by LI Gilbert, K Iatrou & SS Gill), pp. 53–105. Elsevier, Oxford, UK.
- Kilin D, Nagata T & Masuda T (1981) Development of carbamate resistance in the brown planthopper, *Nilaparvata lugens* (Stål). *Applied Entomology and Zoology* 16: 1–6.
- Lan YQ & Zhao SX (2001) The stability of resistance to three pyrethroids in *Spodoptera exigua* Hübner. *Chinese Journal of Pesticide Science* 6: 77–80.
- Le Goff G, Boundy S, Daborn PJ, Yen JL, Sofer L et al. (2003) Microarray analysis of cytochrome P450 mediated insecticide resistance in *Drosophila*. *Insect Biochemistry and Molecular Biology* 33: 701–708.
- LeOra Software (1997) POLO-PC: Probit and Logit Analysis. Leora Software, Berkeley, CA, USA.
- Liu ZW & Han ZJ (2006) Fitness costs of laboratory-selected imidacloprid resistance in the brown planthopper, *Nilaparvata lugens* (Stål). *Pest Management Science* 62: 279–282.
- Liu ZW, Han ZJ, Wang YC, Zhang LC, Zhang HW & Liu CJ (2003) Selection for imidacloprid resistance in *Nilaparvata lugens* (Stål): cross-resistance patterns and possible mechanisms. *Pest Management Science* 59: 1355–1359.
- Mohan M & Gujar GT (2003) Local variation in susceptibility of the diamondback moth, *Plutella xylostella* (Linnaeus) to insecticides and role of detoxification enzymes. *Crop Protection* 22: 495–504.
- Mota-Sanchez D, Hollingworth RM, Grafius EJ & Moyer DD (2006) Resistance and cross-resistance to neonicotinoid insecticides and spinosad in the Colorado potato beetle, *Leptinotarsa decemlineata* (Say) (Coleoptera: Chrysomelidae). *Pest Management Science* 62: 30–37.
- Nauen R & Denholm I (2005) Resistance of insect pests to neonicotinoid insecticides: current status and future prospects. *Archives of Insect Biochemistry and Physiology* 58: 200–215.
- Nauen R, Stumpf N & Elbert A (2002) Toxicological and mechanistic studies on neonicotinoid cross resistance in Q-type *Bemisia tabaci* (Hemiptera: Aleyrodidae). *Pest Management Science* 58: 858–875.
- Prabhaker N, Toscano NC, Castle SJ & Henneberry TJ (1997) Selection for imidacloprid resistance in silverleaf whiteflies from the imperial valley and development of a hydroponic bioassay for resistance monitoring. *Pesticide Science* 51: 419–428.
- Qu MJ, Han ZJ, Xu XJ, Shao XL, Tian XZ & Fu ML (2005) Dynamics and risk assessment of triazophos resistance in rice stem borer (*Chilo suppressalis* Walker). *Journal of Nanjing Agriculture University* 28: 38–42.
- Richman DL, Koehler PG & Brenner RJ (1999) Effect of temperature and the synergist piperonyl butoxide on imidacloprid toxicity to the cat flea (Siphonaptera: Pulicidae). *Household and Structural Insects* 92: 1120–1124.
- Sayyed AH, Attique MR, Khaliq A & Wright DJ (2005) Inheritance of resistance and cross-resistance to deltamethrin in *Plutella xylostella* (Lepidoptera: Plutellidae) from Pakistan. *Pest Management Science* 61: 636–642.
- Shen JL & Wu YD (1995) *Insecticide Resistance in Cotton Bollworm and its Management*. China Agricultural Press, pp. 177–280. Beijing, China.
- Stone BF (1962) The inheritance of DDT-resistance in the cattle tick, *Boophilus microplus*. *Australian Journal of Agricultural Research* 13: 984–1007.
- Sun JZ, Fan JC & Xia LR (1996) Studies on the insecticidal activity of imidacloprid and its application in paddy fields against the brown planthopper, *Nilaparvata lugens* (Homoptera: Delphacidae). *Acta Entomologica Sinica* 39: 37–45.
- Tabashnik BE, Groeters FR, Finson N & Johnson MW (1994) Instability of resistance to *Bacillus thuringiensis*. *Biocontrol Science and Technology* 4: 419–426.
- Tang ZH, Tao LM & Li Z (2006) Resistance of insect pests to neonicotinoid insecticides and management strategies. *Chinese Journal of Pesticide Science* 8: 195–202.
- Wang SC, Ku TY & Chu YI (1988) Resistance patterns in the brown planthopper *Nilaparvata lugens* (Homoptera: Delphacidae) after selection with four insecticides and their combinations. *Plant Protection Bulletin* 30: 59–67.

- Wang YH & Wang MH (2007) Factors affecting the outbreak and management tactics of brown planthopper, *Nilaparvata lugens* (Stål) in China in recent years. *Pesticide Science and Administration* 29: 49–54.
- Wang YH, Shen JL & Wang MH (2005) Research advance on resistance mechanism and management of *Nilaparvata lugens* (Stål) to chemical insecticides. *Pesticide Science and Administration* 26: 24–28.
- Wen ZM & Scott JG (1997) Cross-resistance to imidacloprid in strains of German cockroach (*Blattella germanica*) and housefly (*Musca domestica*). *Pesticide Science* 49: 367–371.
- Wu YD, Shen JL, Tan FJ & You ZP (1996) Stability of pyrethroids resistance in *Helicoverpa armigera* (Hübner). *Acta Entomologica Sinica* 39: 342–346.
- Yu SJ & McCord EJ (2007) Lack of cross-resistance to indoxacarb in insecticide-resistant *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and *Plutella xylostella* (Lepidoptera: Yponomeutidae). *Pest Management Science* 63: 63–67.
- Zhao JZ, Zhu GR, Xu BY, Ju ZL, Zhu SX & Zhu F (1993) Resistance resuming and cross-resistance of *Plutella xylostella* (L.) in Wuhan, China. *Plant Protection* 6: 13–14.
- Zhao JZ, Bishop BA & Grafius EJ (2000) Inheritance and synergism of resistance to imidacloprid in the Colorado potato beetle (Coleoptera: Chrysomelidae). *Journal of Economic Entomology* 93: 1508–1514.
- Zhao JZ, Collins HL, Li YX, Mau RFL, Thompson GD et al. (2006) Monitoring of diamondback moth (Lepidoptera: Plutellidae) resistance to spinosad, indoxacarb and emamectin benzoate. *Journal of Economic Entomology* 99: 176–181.
- Zhao JY, Liu W, Brown JM & Knowles CO (1995) Insecticide resistance in field and laboratory strains of Western flower thrips (Thysanoptera: Thripidae). *Journal of Economic Entomology* 88: 1164–1170.
- Zhuang YL, Shen JL & Chen Z (1999) The influence of triazophos on the productivity of the different wing-form brown planthopper, *Nilaparvata lugens* (Stål). *Journal of Nanjing Agriculture University* 22: 21–24.
- Zhuang YL, Shen JL, Dai DJ & Zhou WJ (2004) Genetic analysis of resistance to buprofezin in the brown planthopper, *Nilaparvata lugens* (Stål) (Homoptera: Delphacidae). *Acta Entomologica Sinica* 47: 749–753.